REFERENCES

Agarry, O.O, M.T. Olaleye and C.O. Bello-Michael (2005), Comparative **REFERENCES**

1. Agarry, O.O, M.T. Olaleye and C.O. Bello-Michael (2005), Comparative antimicrobial activities of *Aloe vera* gel and leaf, African Journal of Biotechnology, 4(12).,1413-1414.
2. Aggarwal D, Bama KS (2004), Tissue culture propagation of elite plant of Aloe vera, L., J Plant BiochemBiotechnol,13.,77–79.
3. Amoo SO, Aremu AO, Van Staden J (2012), In vitro plant regeneration, secondary metabolite production and antioxidant activity of micropropagated Aloe arborescens Mill., Plant Cell Tiss Organ Cult., 11,345–358.
4. Amoo SO, Aremu AO, Van Staden J (2013),Shoot proliferation and rooting treatments influence secondary metabolite production and antioxidant activity in tissue culture-derived Aloe arborescens grown ex vitro, Plant Growth Regul, 70.,115–122.
5. Arnholdt-Schmitt B, Schaffer S (2001), Characterization of genome variation in tissue cultures by RAPD fingerprinting–a methodological comment, Plant Biosyst, 135.,115–120.
6. Awawdeh L, Al-Beitawi M, Hammad M (2009). Effectiveness of propolis and calcium hydroxide as a short-term intracanal medicament against Enterococcus faecalis: a laboratory study. Aust. Endod. J, 35., 52-58.
7. Bantawa P, Saha-Roy O, Kumar S, Ghosh R, Mondal K.T. (2011),*In vitro* regeneration of an endangered medicinal plant,*Picrorhizscrophulariiflora*. Biol. Plant,55.,169–172.
8. Bedini C, Caccia R, Triggiani D, Mazzucato A, Soressi GP, Tiezzi A (2009) , Micropropagation of Aloe arborescens Mill: a step towards efficient production of its

valuable leaf extracts showing anti proliferative activity on murine myeloma cells, Plant Biosyst,143(2).,233–240.

9. Campestrini LH, Kuhnen S, Lemos PMM, Bach DB, Dias PF and Maraschin M (2006). Cloning protocol of Aloe vera as a study-case for tailormade biotechnology to small farmers. J TechnolManagInnov, 1.,76-79
10. Cavallini, A, Natali, L, Cionini, C, Sessoli, O. and Sanchez, I.C. (1993),*Invitro* culture of *Aloe barbadensis* Mill. Quantitative DNA variation in regenerated plants, Plant Science,91.,223-224.
11. Chaudhary AK, Ray AK, Jha S, Mishra IN (2011), Callus formation, shoot initiation and in vitro culture of *Aloe vera*, Biotechnology, Bioinf. Bioeng, 1 (4).,551-553.
12. Cheruvathur MK, Abraham J, Thomas TD (2013), Plant regeneration through callus organogenesis and true-to-type conformity of plants by RAPD analysis in Desmodiumgangeticum (Linn.) DC. ApplBiochemBiotechnol,169.,1799–1810.
13. CHUKWUJEKWU J.C., FENNELL C.W., VAN STADEN J. 2002. Optimization of the tissue culture protocol for the endangered Aloe polyphylla. South African Journal of Botany, 68.,424-429.
14. Cragg GM, Newman DJ, Weiss RB. Coral Reefs, Forests, and Thermals Vents: The Worldwide Exploration of Nature for Novel Antitumor Agents. SeminOncol 1997, 24.,156-63.
15. Davis, R.H. and Leiter, M.G. (1988), Aloe Vera: A natural approach for treating wounds, edema, and pain diabetes, Journal of The American Podiatric Medical Association, 78(2).,60-68.
16. Debiasi C, Silva CG, PescadorR. Plant. Med. Botucatu. 2007, 9.,36-43.
17. De Oliveira ET, Crocomo OJ (2009), Large-scale micropropagation of Aloe vera. HortSci 44(6).,1675–1678.

18. Eshun K, He Q (2004), Aloe vera: a valuable ingredient for the food, pharmaceutical and cosmetic industries–a review. Crit Rev Food SciNutr, 44.,91–96
19. Farnsworth NR, Akerele O, Bingel AS, Soejarto DD, Guo Z. Medicinal Plants in Therapy. Bull World Health Organ 1985, 63.,965-81.
20. Gadzovska-Simic,S.(2010)Secondary metabolite production in cultures *in vitro* of *Hypericum perforatum* L.Effects of different exogenous factors, Lambert Academic Publishing, Saarbrücken,Germany, pp., 1-192.
21. GamborgO.L,MillerR.A,andK.Ojima,(1968), Nutrient requirements of suspension cultures soybean root cells, *Exp. Cell Res*.**50.**,148-151.
22. Gantait S, Mandal N, Das PK (2011), In vitro accelerated mass propagation and ex vitro evaluation of Aloe vera L with aloin content and superoxide dismutase activity, Nat Prod Resh,25(14).,1370–1378.
23. Goswami K, Sharma R, Singh PK, Singh G (2013) Micropropagation of seedless lemon (Citrus limon L. cv. KaghziKalan) and assessment of genetic fidelity of micropropagated plants using RAPD markers. PhysiolMolBiol Plants, 19., 137–145
24. Grace OM, Simmon MSJ, Smith GF, Van Wyk AE (2008), Therapeutic uses of Aloe L (Asphodelaceae) in Southern Africa, J Ethnopham ,119.,604–614.
25. Harirah AA, Khalid N,(2006), Direct regeneration and RAPD assessment of male inflorescence derived plants of Musa acuminata Cv. Berangan, Asia Pac J MolBiol Biotechnology, 14., 11–17.
26. Hashemabadi D, Kaviani B, (2008), Rapid micropropagation of Aloe vera L. via shoot multiplication, African Jour. Biotechnology, 7 (12).,1899-1902.
27. Hashemabadi D, Kaviani B , (2008), In vitro proliferation of an important medicinal plant Aloe- A method for rapid production, Australian Journal of Crop Science, 4 (4)., 216-222.

28. Haque SM, Ghosh B (2013a), Field evaluation and genetic stability assessment of regenerated plants produced via direct shoot organogenesis from leaf explant of an endangered ‘Asthma Plant’ (Tylophoraindica) along with their in vitro conservation, NatlAcadSciLett, doi 10, 1007/s40009-013-0161-z.
29. Haque SM, Ghosh B (2013b), Micropropagation, in vitro flowering and cytological studies of Bacopachamaedryoides, an ethno-medicinal plant. EnvExpBiol, 11.,59–68.
30. HOSSEINI R., PARSA M. 2002. Micropropagation of Aloe vera L. grown in South Iran. Pakistan Journal of Biological Sciences 10.,1134-1137.
31. Hayacibara MF, Koo H, Rosalen PL, Duarte S, Franco EM, Bowen WH, Ikegaki M, Cury JA (2005). In vitro and in vivo effects of isolated fractions of Brazilian propolis on caries development. J. Ethnopharmacol., 101.,110-115.
32. Jayakrishna C, Karthik, Barathi S, Kamalanathan D, IndraArulselvi P (2011), *In vitro* propagation of *Aloe barbadensis* Miller, a miracle herb, Research in Plant Biology, 1 (5).,22-26.
33. Joshi, P. and V. Dhawan, 2007.Assessment of genetic fidelity of micropropagatedSwertiachirayita plantlets by ISSR marker assay. Biol. Plantarum, 51.,22-26.
34. Kametani, S. Yuasa, A.K. Kikuzaki, H. Kennedy, D.O. Honzawa, M. Yuasa, M. (2007), Chemical constituents of Cape Aloe and their synergistic growth inhibiting effect on Ehrlich Ascites Tumor Cells, Biosci. Biotechnol. Biochem, 71(5).,1220-1229.
35. Kathi JK, Victoria C (1999), Aloe vera. The Longwood Herbal Task Force and the Centre for Holistic Pediatric Education and Research, 2.,29-34.

36. Kumar A, Prakash K, Sinha RK, Kumar N (2013), In vitro plant propagation of Catharanthusroseus and assessment of genetic fidelity of micropropagated plants by RAPD marker assay, ApplBiochem Biotech 169., 894–900.
37. Kumi-Diaka J, Nguyen V, Butter A. Cytotoxic potential of the phytochemical genisteinisoflavone and certain environmental chemical compound on testicular cells, Biol cell 1999, 91., 515-23.
38. Lad VN, Murthy ZVP (2013), Rheology of Aloe barbadensis Miller: a naturally available material of high therapeutic and nutrient value for food applications, J Food Eng ,115.,279–284.
39. Langmead L, Chitnis M, Rampton DS. Use of complementarytherapies by patients with IBD may indicate psychosocialdistress. Inflamm Bowel Dis 2002, 81.,74–9.
40. Lawless, J. Allan, J. (2000), The Clinical Composition of Aloe vera, InAloeveranatural wonder cure. Thorsons, Publishing Ltd., London, United Kingdom. Pp, 161-171.
41. Leelambika M, Sathyanarayana N (2011), Genetic characterization of Indian Mucuna (Leguminoceae) species using morphometric and RAPD approaches, Plant Biosyst, 145(4)., 786–797.
42. Liao Z, Chen M, Tan F, Sun X, Tang K (2004),Micropropagation of endangered Chinese aloe. Plant Cell, Tiss. Organ Cult,78.,83-86.
43. Martins M, Sarmento D, Oliveira MM (2004), Genetic stability of micropropagated almond plantlets, as assessed by RAPD and ISSR markers, Plant Cell Rep, 23.,492–496.
44. Meyer HJ, Staden J (1991), Rapid in vitro propagation of *Aloe barhadensis* Mill. Plant Cell Tiss. Org. Cult, 76 (1).,83-86.

45. Mohanty S, Panda M.K, Sahoo S. &Nayak S. (2011),Micropropagation of Zingiberrubens and assessment of genetic stability through RAPD and ISSR markers, Biol. Plant, 55., 16–20.
46. MulinacciN,GiaccheriniC,SantamariaA,CaniatoR,FerrariF,Valletta A,VincieriF,andG.Pasqua(2008),Anthocyaninsandxanthonesinthecalliandregeneratedshootsof*Hypericumperforatum*var.*angustifolium*(sin. Fröhlich)Borkh,*Plant*Physiol.Biochem,46.,414-420.
47. Natali L, Sanchez IC, Cavallini (1990), In vitro culture of *Aloe barbadensis* Mill. micropropagation from vegetative meristems, Plant Cell, Tiss. Org. cult,20.,71-74.
48. Ni Y, Turner D, Yates K.M,Tizard,(2004) , Isolation and characterisation of structural components of *Aloe vera*L, leaf pulp,Int. Immunopharmaco,*4.,* 1745-1755.
49. NiY,Yates K.M, Tizard, I.R(2004), Aloe polysaccharides, In *Aloes The Genus Aloe;* Reynolds, T,Ed, CRC Press: Boca Raton, pp, 75-87.
50. Oncag O, Cogulu D, Uzel A, Sorkun K (2008). Efficacy of propolis as an intracanal medicament against Enterococcus faecalis. General Dentistry, 54(5).,319-322.
51. Ozan F, Sümer Z, Polat ZA, Er K, Ozan U, Deer O (2007). Effect of mouth rinse containing propolis on oral microorganisms and human gingival fibroblast. Eur. J. dentistry, 11.,195-200.
52. Pasqua,G., Avato,P., Monacelli,B., Santamaria,A., and M.Argentieri(2003) Metabolites in cellsus pensioncultures, calli, and *in vit*ro regenerated organs of *Hypericumperforatum*cv. Topas, PlantSci, 16., 977-982.
53. Paridaa R, Mohantya S, Nayak S (2013)*In vitro* propagation of *Hedychiumcoronarium*Koen: through axillary bud proliferation, Plant Biosyst, doi:10.1080/11263504.2012.748102.
54. Premvaranon P, Vearasilp S, Thanapornpoong S, KarladeeD, and Gorinstein S, (2011) *In vitro* studies to produce double haploid inIndica hybrid rice,Biologia, 66., 1074–

1081.

55. Rahman M, Rajora O,(2001), Microsatellite DNA somaclonal variations in micropropagated trembling aspen (*Populustremuloides*), Plant Cell Rep, 20., 531–536.
56. Rathore MS, Chikara J, Mastan SG, Rahman H, Anand KGV, Shekhawat NS (2011a), Assessment of genetic stability and instability of tissue culture-propagated plantlets of Aloe vera L by RAPD and ISSR Markers,Applied Biochem Biotechnol, 165.,1356–1365.
57. Rout GR, Das P, Goel S, Raina SN, (1998), Determination of genetic stability of micropropagated plants of ginger using random amplified polymorphic DNA (RAPD) markers, Bot Bull Acad Sin, 39.,23–27.
58. Rout GR, (2002), Direct plant regeneration from leaf explants of plumbago species and its genetic fidelity through RAPD markers, Ann ApplBiol, 140., 305–313.
59. Rout G.R, Mahato A, Senapati S.K, (2008), *In vitro* clonal propagation of *Nyctanthesarbortristis*, Biol. Plant, 52.,521– 524.
60. Roy S.C, and Sarkar A, (1991), In vitro regeneration and micro propagation of *Aloe vera*, Scientia Horticulturae, 47(1-2)., 107-114.
61. Sabir A, Tabbu CR, Agustiono P, Sosroseno W (2005). Histological analysis of rat dental pulp tissue capped with propolis. J. Oral Sci., 47(3).,135-138.
62. Samantaray S, Maiti S (2008), Rapid plant regeneration and assessment of genetic fidelity of in vitro raised plants in Aloe barbadensis Mill: using RAPD markers, Acta Bot Gallica 155(3)., 427–434.
63. Sanche I.C, Natali L, and Cavallini, A, (1988), In vitro culture of *Aloe barbadensis* Mill. Morphogenetic ability and nuclear DNA content, Plant Science, 55.,53-59.

64. Savita, Bhagat A, Pati PK, Virk GS, Nagpal A (2012),An efficient micropropagation protocol for Citrus jambhiri Lush. and assessment of clonal fidelity employing anatomical studies and RAPD markers, In Vitro Cell &DevBiol–Plant,48.,512–520.

65. Senapati S, Aparajita S, Rout GR (2013),Micropropagation and assessment of genetic stability in *Celastruspaniculatus*: An endangered medicinal plant, Biologia, 68/4., 627—632.

66. Shukla MR, Jones AMP, Sullivan JA, Liu CZ, Gosling S, Saxena PK (2012), In vitro conservation of American elm (Ulmusamericana): potential role of auxin metabolism in sustained plant proliferation, Can J Forest Re,42.,686–697.

67. Singh M, Rathore MS, Panwar D, Rathore JS, Dagla HR, Shekhawat NS (2009),Micropropagation of selected genotype of Aloe vera L–an ancient plant for modern industry, J Sustain Forest, 28(8).,935–950.

68. Swarna J., Ravindhran R. (2012), In vitro propagation and assessment of genetic integrity of *Talinumtriangulare* (Jacq.) Wild: a valuable medicinal herb, ActaPhysiol Plant, 10., 1007–1017.

69. Talmadge, J. Chavez, J. Jacobs, L.Munger, C. Chinnah, T.Chow, J.T. Williamson, D. Yates,K.(2004), Fractionation of *Aloe vera*L. inner gel, purification and molecular profiling of activity. *Int.Immunopharmacol,4.,* 1757-1773.

70. Tan, B. K., & Vanitha, J. (2004). Immunomodulatory and antimicrobial effects of some traditional Chinese medicinal herbs: a review. Current Medicinal Chemistry, 11(11)., 1423–1430.

71. Tanaka M, Misawa E, Ito Y, Habara N, Nomaqushi K, Yamada M, Toida T, Hayasawa H, Takase M, Inaqaki MandHiqushi R. Identification of five phytosterols from Aloe vera gel as anti-diabetic compounds. Biol. Pharm. Bull. 2006, 29 (7)., 1418-22.

72. Toker H, Ozan F, Ozer H, Ozdemir H, Eren K, Yeler HJ (2008). A morphometic and histopathologic evaluation of the effects of Propolis on alveolar bone loss in experimental periodontitis in rats. Periodontol, 79(6)., 1089-1094.
73. Vogler BK, Ernst E (1999)*Aloe vera*: a systematic review of its clinical effectiveness, Brit J Gen Pract,49.,823–828.
74. Yagi A, Kabbash A, Mizuno K, Moustafa SM, Khalifa TI and Tsuji H. Radical scavenging glycoprotein inhibiting cyclooxygenase-2 and thromboxane A2 synthase from Aloe vera gel. Planta Med, 2006, 69.,69-71.
75. Yongchaiyudha S, Rungpitarangsi V, Bunyapraphatsara N, and Chokechaijaroenporn O. (1996) Antidiabetic activity of Aloe vera juice l. Clinical trial in new cases of diabetes mellitus. Phytomedicine, 3., 241-43.
76. Yuan, Y,J, Li,J.C, HuZ.D, WuJ.C, and Zeng, A.P. (2002), Fungal elicitor-induced cell apoptosis in suspension cultures of *Taxus chinensis* var. Maireifortaxol production. Proc.Biochem,38.,193-198.
77. Zapata PJ, Navarro D, GuillénF, Castillo S, Martínez-Romero D, Valero D, Serrano M (2013), Characterization of gels from different Aloe spp as antifungal treatment: Potential crops for industrial applications, Ind Crop Prod,42.,223–230.
78. Zhao J, Davis L, and Verpoorte, R(2005) Elicitor signal transduction leading to production of plant secondary metabolites. Biotechnol. Adv,23.,283–333.
79. antimicrobial activities of *Aloe vera* gel and leaf, African Journal of Biotechnology, 4(12).,1413-1414.
80. Aggarwal D, Bama KS (2004), Tissue culture propagation of elite plant of Aloe vera, L., J Plant BiochemBiotechnol,13.,77–79.

81. Amoo SO, Aremu AO, Van Staden J (2012), In vitro plant regeneration, secondary metabolite production and antioxidant activity of micropropagated Aloe arborescens Mill., Plant Cell Tiss Organ Cult., 11,345–358.
82. Amoo SO, Aremu AO, Van Staden J (2013),Shoot proliferation and rooting treatments influence secondary metabolite production and antioxidant activity in tissue culture-derived Aloe arborescens grown ex vitro, Plant Growth Regul, 70.,115–122.
83. Arnholdt-Schmitt B, Schaffer S (2001), Characterization of genome variation in tissue cultures by RAPD fingerprinting–a methodological comment, Plant Biosyst, 135.,115–120.
84. Awawdeh L, Al-Beitawi M, Hammad M (2009). Effectiveness of propolis and calcium hydroxide as a short-term intracanal medicament against Enterococcus faecalis: a laboratory study. Aust. Endod. J, 35., 52-58.
85. Bantawa P, Saha-Roy O, Kumar S, Ghosh R, Mondal K.T. (2011),*In vitro* regeneration of an endangered medicinal plant,*Picrorhizscrophulariiflora*. Biol. Plant,55.,169–172.
86. Bedini C, Caccia R, Triggiani D, Mazzucato A, Soressi GP, Tiezzi A (2009) , Micropropagation of Aloe arborescens Mill: a step towards efficient production of its valuable leaf extracts showing anti proliferative activity on murine myeloma cells, Plant Biosyst,143(2).,233–240.
87. Campestrini LH, Kuhnen S, Lemos PMM, Bach DB, Dias PF and Maraschin M (2006). Cloning protocol of Aloe vera as a study-case for tailormade biotechnology to small farmers. J TechnolManagInnov, 1.,76-79
88. Cavallini, A, Natali, L, Cionini, C, Sessoli, O. and Sanchez, I.C. (1993),*Invitro* culture of *Aloe barbadensis* Mill. Quantitative DNA variation in regenerated plants, Plant

Science,91.,223-224.

89. Chaudhary AK, Ray AK, Jha S, Mishra IN (2011), Callus formation, shoot initiation and in vitro culture of *Aloe vera*, Biotechnology, Bioinf. Bioeng, 1 (4).,551-553.
90. Cheruvathur MK, Abraham J, Thomas TD (2013), Plant regeneration through callus organogenesis and true-to-type conformity of plants by RAPD analysis in Desmodiumgangeticum (Linn.) DC. ApplBiochemBiotechnol,169.,1799–1810.
91. CHUKWUJEKWU J.C., FENNELL C.W., VAN STADEN J. 2002. Optimization of the tissue culture protocol for the endangered Aloe polyphylla. South African Journal of Botany, 68.,424-429.
92. Cragg GM, Newman DJ, Weiss RB. Coral Reefs, Forests, and Thermals Vents: The Worldwide Exploration of Nature for Novel Antitumor Agents. SeminOncol 1997, 24.,156-63.
93. Davis, R.H. and Leiter, M.G. (1988), Aloe Vera: A natural approach for treating wounds, edema, and pain diabetes, Journal of The American Podiatric Medical Association, 78(2).,60-68.
94. Debiasi C, Silva CG, PescadorR. Plant. Med. Botucatu. 2007, 9.,36-43.
95. De Oliveira ET, Crocomo OJ (2009), Large-scale micropropagation of Aloe vera. HortSci 44(6).,1675–1678.
96. Eshun K, He Q (2004), Aloe vera: a valuable ingredient for the food, pharmaceutical and cosmetic industries–a review. Crit Rev Food SciNutr, 44.,91–96
97. Farnsworth NR, Akerele O, Bingel AS, Soejarto DD, Guo Z. Medicinal Plants in Therapy. Bull World Health Organ 1985, 63.,965-81.
98. Gadzovska-Simic,S.(2010)Secondary metabolite production in cultures *in vitro* of *Hypericum perforatum* L.Effects of different exogenous factors, Lambert Academic Publishing, Saarbrücken,Germany, pp., 1-192.

99. GamborgO.L,MillerR.A,andK.Ojima,(1968), Nutrient requirements of suspension cultures soybean root cells, *Exp. Cell Res*.**50.**,148-151.

100. Gantait S, Mandal N, Das PK (2011), In vitro accelerated mass propagation and ex vitro evaluation of Aloe vera L with aloin content and superoxide dismutase activity, Nat Prod Resh,25(14).,1370–1378.

101. Goswami K, Sharma R, Singh PK, Singh G (2013) Micropropagation of seedless lemon (Citrus limon L. cv. KaghziKalan) and assessment of genetic fidelity of micropropagated plants using RAPD markers. PhysiolMolBiol Plants, 19., 137–145

102. Grace OM, Simmon MSJ, Smith GF, Van Wyk AE (2008), Therapeutic uses of Aloe L (Asphodelaceae) in Southern Africa, J Ethnopham ,119.,604–614.

103. Harirah AA, Khalid N,(2006), Direct regeneration and RAPD assessment of male inflorescence derived plants of Musa acuminata Cv. Berangan, Asia Pac J MolBiol Biotechnology, 14., 11–17.

104. Hashemabadi D, Kaviani B, (2008), Rapid micropropagation of Aloe vera L. via shoot multiplication, African Jour. Biotechnology, 7 (12).,1899-1902.

105. Hashemabadi D, Kaviani B , (2008), In vitro proliferation of an important medicinal plant Aloe- A method for rapid production, Australian Journal of Crop Science, 4 (4)., 216-222.

106. Haque SM, Ghosh B (2013a), Field evaluation and genetic stability assessment of regenerated plants produced via direct shoot organogenesis from leaf explant of an endangered ‘Asthma Plant’ (Tylophoraindica) along with their in vitro conservation, NatlAcadSciLett, doi 10, 1007/s40009-013-0161-z.

107. Haque SM, Ghosh B (2013b), Micropropagation, in vitro flowering and cytological studies of Bacopachamaedryoides, an ethno-medicinal plant. EnvExpBiol, 11.,59–68.

108. HOSSEINI R., PARSA M. 2002. Micropropagation of Aloe vera L. grown in South Iran. Pakistan Journal of Biological Sciences 10.,1134-1137.

109. Hayacibara MF, Koo H, Rosalen PL, Duarte S, Franco EM, Bowen WH, Ikegaki M, Cury JA (2005). In vitro and in vivo effects of isolated fractions of Brazilian propolis on caries development. J. Ethnopharmacol., 101.,110-115.

110. Jayakrishna C, Karthik, Barathi S, Kamalanathan D, IndraArulselvi P (2011), *In vitro* propagation of *Aloe barbadensis* Miller, a miracle herb, Research in Plant Biology, 1 (5).,22-26.

111. Joshi, P. and V. Dhawan, 2007.Assessment of genetic fidelity of micropropagatedSwertiachirayita plantlets by ISSR marker assay. Biol. Plantarum, 51.,22-26.

112. Kametani, S. Yuasa, A.K. Kikuzaki, H. Kennedy, D.O. Honzawa, M. Yuasa, M. (2007), Chemical constituents of Cape Aloe and their synergistic growth inhibiting effect on Ehrlich Ascites Tumor Cells, Biosci. Biotechnol. Biochem, 71(5).,1220-1229.

113. Kathi JK, Victoria C (1999), Aloe vera. The Longwood Herbal Task Force and the Centre for Holistic Pediatric Education and Research, 2.,29-34.

114. Kumar A, Prakash K, Sinha RK, Kumar N (2013), In vitro plant propagation of Catharanthusroseus and assessment of genetic fidelity of micropropagated plants by RAPD marker assay, ApplBiochem Biotech 169., 894–900.

115. Kumi-Diaka J, Nguyen V, Butter A. Cytotoxic potential of the phytochemical genisteinisoflavone and certain environmental chemical compound on testicular cells, Biol cell 1999, 91., 515-23.

116. Lad VN, Murthy ZVP (2013), Rheology of Aloe barbadensis Miller: a naturally available material of high therapeutic and nutrient value for food applications, J Food Eng ,115.,279–284.

117. Langmead L, Chitnis M, Rampton DS. Use of complementarytherapies by patients with IBD may indicate psychosocialdistress. Inflamm Bowel Dis 2002, 81.,74–9.

118. Lawless, J. Allan, J. (2000), The Clinical Composition of Aloe vera, InAloeveranatural wonder cure. Thorsons, Publishing Ltd., London, United Kingdom. Pp, 161-171.

119. Leelambika M, Sathyanarayana N (2011), Genetic characterization of Indian Mucuna (Leguminoceae) species using morphometric and RAPD approaches, Plant Biosyst, 145(4)., 786–797.

120. Liao Z, Chen M, Tan F, Sun X, Tang K (2004),Micropropagation of endangered Chinese aloe. Plant Cell, Tiss. Organ Cult,78.,83-86.

121. Martins M, Sarmento D, Oliveira MM (2004), Genetic stability of micropropagated almond plantlets, as assessed by RAPD and ISSR markers, Plant Cell Rep, 23.,492–496.

122. Meyer HJ, Staden J (1991), Rapid in vitro propagation of *Aloe barhadensis* Mill. Plant Cell Tiss. Org. Cult, 76 (1).,83-86.

123. Mohanty S, Panda M.K, Sahoo S. &Nayak S. (2011),Micropropagation of Zingiberrubens and assessment of genetic stability through RAPD and ISSR markers, Biol. Plant, 55., 16–20.

124. MulinacciN,GiaccheriniC,SantamariaA,CaniatoR,FerrariF,VallettaA,VincieriF,andG.Pasqua(2008),Anthocyaninsandxanthonesinthecalliandregeneratedshootsof*Hypericumperforatum*var.*angustifolium*(sin. Fröhlich)Borkh,*Plant*Physiol.Biochem,46.,414-420.

125. Natali L, Sanchez IC, Cavallini (1990), In vitro culture of *Aloe barbadensis* Mill. micropropagation from vegetative meristems, Plant Cell, Tiss. Org. cult,20.,71-74.

126. Ni Y, Turner D, Yates K.M,Tizard,(2004) , Isolation and characterisation of structural components of *Aloe vera*L, leaf pulp,Int. Immunopharmaco,*4.*, 1745-1755.

127. NiY,Yates K.M, Tizard, I.R(2004), Aloe polysaccharides, In *Aloes The Genus Aloe;* Reynolds, T,Ed, CRC Press: Boca Raton, pp, 75-87.

128. Oncag O, Cogulu D, Uzel A, Sorkun K (2008). Efficacy of propolis as an intracanal medicament against Enterococcus faecalis. General Dentistry, 54(5).,319-322.

129. Ozan F, Sümer Z, Polat ZA, Er K, Ozan U, Deer O (2007). Effect of mouth rinse containing propolis on oral microorganisms and human gingival fibroblast. Eur. J. dentistry, 11.,195-200.

130. Pasqua,G., Avato,P., Monacelli,B., Santamaria,A., and M.Argentieri(2003) Metabolites in cellsus pensioncultures, calli, and *in vit*ro regenerated organs of *Hypericumperfo- ratum*cv. Topas, PlantSci, 16., 977-982.

131. Paridaa R, Mohantya S, Nayak S (2013)*In vitro* propagation of *Hedychiumcoronarium*Koen: through axillary bud proliferation, Plant Biosyst, doi:10.1080/11263504.2012.748102.

132. Premvaranon P, Vearasilp S, Thanapornpoong S, KarladeeD, and Gorinstein S, (2011) *In vitro* studies to produce double haploid inIndica hybrid rice,Biologia, 66.,

1074–1081.

133. Rahman M, Rajora O,(2001), Microsatellite DNA somaclonal variations in micropropagated trembling aspen (*Populustremuloides*), Plant Cell Rep, 20., 531–536.

134. Rathore MS, Chikara J, Mastan SG, Rahman H, Anand KGV, Shekhawat NS (2011a), Assessment of genetic stability and instability of tissue culture-propagated plantlets of Aloe vera L by RAPD and ISSR Markers,Applied Biochem Biotechnol, 165.,1356–1365.

135. Rout GR, Das P, Goel S, Raina SN, (1998), Determination of genetic stability of micropropagated plants of ginger using random amplified polymorphic DNA (RAPD) markers, Bot Bull Acad Sin, 39.,23–27.

136. Rout GR, (2002), Direct plant regeneration from leaf explants of plumbago species and its genetic fidelity through RAPD markers, Ann ApplBiol, 140., 305–313.

137. Rout G.R, Mahato A, Senapati S.K, (2008), *In vitro* clonal propagation of *Nyctanthesarbortristis*, Biol. Plant, 52.,521– 524.

138. Roy S.C, and Sarkar A, (1991), In vitro regeneration and micro propagation of *Aloe vera*, Scientia Horticulturae, 47(1-2)., 107-114.

139. Sabir A, Tabbu CR, Agustiono P, Sosroseno W (2005). Histological analysis of rat dental pulp tissue capped with propolis. J. Oral Sci., 47(3).,135-138.

140. Samantaray S, Maiti S (2008), Rapid plant regeneration and assessment of genetic fidelity of in vitro raised plants in Aloe barbadensis Mill: using RAPD markers, Acta Bot Gallica 155(3)., 427–434.

141. Sanche I.C, Natali L, and Cavallini, A, (1988), In vitro culture of *Aloe barbadensis* Mill. Morphogenetic ability and nuclear DNA content, Plant Science, 55.,53-59.

142. Savita, Bhagat A, Pati PK, Virk GS, Nagpal A (2012),An efficient micropropagation protocol for Citrus jambhiri Lush. and assessment of clonal fidelity employing anatomical studies and RAPD markers, In Vitro Cell &DevBiol–Plant,48.,512–520.

143. Senapati S, Aparajita S, Rout GR (2013),Micropropagation and assessment of genetic stability in *Celastruspaniculatus*: An endangered medicinal plant, Biologia, 68/4., 627—632.

144. Shukla MR, Jones AMP, Sullivan JA, Liu CZ, Gosling S, Saxena PK (2012), In vitro conservation of American elm (Ulmusamericana): potential role of auxin metabolism in sustained plant proliferation, Can J Forest Re,42.,686–697.

145. Singh M, Rathore MS, Panwar D, Rathore JS, Dagla HR, Shekhawat NS (2009),Micropropagation of selected genotype of Aloe vera L–an ancient plant for modern industry, J Sustain Forest, 28(8).,935–950.

146. Swarna J., Ravindhran R. (2012), In vitro propagation and assessment of genetic integrity of *Talinumtriangulare* (Jacq.) Wild: a valuable medicinal herb, ActaPhysiol Plant, 10., 1007–1017.

147. Talmadge, J. Chavez, J. Jacobs, L.Munger, C. Chinnah, T.Chow, J.T. Williamson, D. Yates,K.(2004), Fractionation of *Aloe vera*L. inner gel, purification and molecular profiling of activity. *Int.Immunopharmacol,4.*, 1757-1773.

148. Tan, B. K., & Vanitha, J. (2004). Immunomodulatory and antimicrobial effects of some traditional Chinese medicinal herbs: a review. Current Medicinal Chemistry, 11(11)., 1423–1430.

149. Tanaka M, Misawa E, Ito Y, Habara N, Nomaqushi K, Yamada M, Toida T, Hayasawa H, Takase M, Inaqaki MandHiqushi R. Identification of five phytosterols

from Aloe vera gel as anti-diabetic compounds. Biol. Pharm. Bull. 2006, 29 (7)., 1418-22.

150. Toker H, Ozan F, Ozer H, Ozdemir H, Eren K, Yeler HJ (2008). A morphometic and histopathologic evaluation of the effects of Propolis on alveolar bone loss in experimental periodontitis in rats. Periodontol, 79(6)., 1089-1094.

151. Vogler BK, Ernst E (1999)*Aloe vera*: a systematic review of its clinical effectiveness, Brit J Gen Pract,49.,823–828.

152. Yagi A, Kabbash A, Mizuno K, Moustafa SM, Khalifa TI and Tsuji H. Radical scavenging glycoprotein inhibiting cyclooxygenase-2 and thromboxane A2 synthase from Aloe vera gel. Planta Med, 2006, 69.,69-71.

153. Yongchaiyudha S, Rungpitarangsi V, Bunyapraphatsara N, and Chokechaijaroenporn O. (1996) Antidiabetic activity of Aloe vera juice l. Clinical trial in new cases of diabetes mellitus. Phytomedicine, 3., 241-43.

154. Yuan, Y,J, Li,J.C, HuZ.D, WuJ.C, and Zeng, A.P. (2002), Fungal elicitor-induced cell apoptosis in suspension cultures of *Taxus chinensis* var. Maireifortaxol production. Proc.Biochem,38.,193-198.

155. Zapata PJ, Navarro D, GuillénF, Castillo S, Martínez-Romero D, Valero D, Serrano M (2013), Characterization of gels from different Aloe spp as antifungal treatment: Potential crops for industrial applications, Ind Crop Prod,42.,223–230.

156. Zhao J, Davis L, and Verpoorte, R(2005) Elicitor signal transduction leading to production of plant secondary metabolites. Biotechnol. Adv,23.,283–333.